

Extracorporeal membrane oxygenation in neonates and children

Martin Schmiady^a, Carsten Döll^b, Anna Cavigelli-Brunner^c, Alexander Konetzka^a, Sebastian Bründler^b, Yvonne Kröger^b, Hitendu Dave^a, Michael Hübler^a, Martin Schweiger^a

^a University Children's Hospital Zurich, Division of Congenital Cardiovascular Surgery, Switzerland

^b University Children's Hospital Zurich, Department of Paediatric Intensive Care and Neonatology, Switzerland

^c University Children's Hospital Zurich, Division of Paediatric Cardiology, Switzerland

Summary

Extracorporeal membrane oxygenation (ECMO) is a general term that describes the short- or long-term support of the heart and/or lungs in neonates, children and adults. It offers a treatment option for severe cardiac and/or respiratory failure from neonate to adult. Used as venoarterial extracorporeal membrane oxygenation (VA-ECMO), it remains the most commonly used modality for short- to mid-term mechanical support of the failing circulation in children. The aim of this article is to review the clinical indications, different circuits, technical options, patient management, and potential risks and benefits of this therapy for children. As ECMO therapy is an overwhelming event for the whole family we also highlight the role of psychosocial counselling and support for the parents.

Key words: ECMO; neonates; children; cannulation, management

ity to provide support for both lung and heart function is the major difference of VA-ECMO from VV-ECMO. In VV-ECMO the artificial lung is situated in series with the native lungs; cardiac function must be adequate to maintain pulmonary and systemic perfusion.

Pulmonary or/and myocardial failure are the main indications for the clinical application of ECMO (table 1). Despite growing experience, some clinical features remain contraindications for ECMO therapy (table 2).

Although patient retrieval on ECMO is feasible, patients with a high probability of requiring ECMO therapy are ideally transferred in good time to ECMO centres with significant expertise in paediatrics [2, 3]. Patient are preferably referred before ongoing deterioration of the underlying medical condition occurs, patient transport is precluded by their condition or secondary additional organ damage is observed [4].



Introduction

Extracorporeal membrane oxygenation (ECMO) is a modified form of cardiopulmonary bypass (CPB) used to provide adequate organ oxygenation in patients with cardiac and/or respiratory failure. The scope of this article is to give general practitioners and specialists not involved in ECMO management a brief overview of indications, surgical implantation and patient/parental management for children treated with ECMO.

Indications and types of ECMO

ECMO is a partial or complete support of ventilation and oxygenation, as well as uni- or biventricular support of myocardial function. It operates by removing blood from the patient's venous circulation, pumping it through an oxygenator to remove carbon dioxide and oxygenate haemoglobin, and subsequently returning it to either the arterial (VA-ECMO) or venous (VV-ECMO) circulation. In VA-ECMO the artificial lung is in parallel with the native lungs. This allows sufficient oxygenation, as well as cardiac output. The abil-

The ECMO circuit

We use a Thoratec Centrimag® or PediVas® (Levitronix, Zurich, Switzerland) console with a back-up unit, a heater unit and a Sechrist Air-Oxygen-Mixer (Sechrist Industries, Anaheim, USA) (fig. 1). For neonates and infants weighing up to 15 kg the whole circuit contains approximately 250 ml priming solution with flow ranges up to 1.7 l/min; for children above 15 kg the volume of priming solution is about 750 ml and flow rates up to 7.0 l/min can be achieved.

Implantation techniques and cannulation sites

Weight limits for different cannulation sites and various cannulation algorithms have been published [5, 6]. Cannulation for neonates, infants and small children focuses on either neck vessels or large central vessels via median sternotomy. Cannulation of the femoral vessels in small children is not possible because the small size of the vessels does not allow implantation of cannulae large enough to achieve full ECMO flow.

Table 1: Extracorporeal membrane oxygenation indications.

Heart failure	Failure to wean from CPB
	Myocarditis
	Cardiomyopathy (bridge to recovery, transplant or long term MCS)
	Refractory sepsis with profound cardiac depression*
	Refractory cardiac arrhythmias
	Coronary ischaemia (if surgically amenable/treatable)
Pulmonary failure	Bridge to transplantation
	Neonatal respiratory distress syndrome
	Adult respiratory distress syndrome [†]
	Persistent pulmonary hypertension of the newborn / persistent fetal circulation
	Meconium aspiration syndrome
	Pneumonia (viral/bacterial/aspiration)
Cardiac arrest from any cause[‡]	Air leak syndrome
	Bridge to decision, underlying treatable disease
Transplantation	Pre-transplantation as bridge to Tx
	Primary graft dysfunction after heart or lung Tx
Elective periprocedural support	During lung transplantation or tracheal surgery
Congenital diaphragmatic hernia	In special circumstances; centre specific

(CPB = cardiopulmonary bypass; MCS = mechanical circulatory support; Tx = transplantation)

* In patients with impaired cardiac output who cannot sufficiently increase cardiac output in sepsis; the team has to be aware that further invasive treatment (surgery, e.g., thoracotomy or laparotomy) on ECMO therapy is associated with higher complication rates (bleeding) than normal [8].

[†] In children and adolescents.

[‡] Depending on clinical situation; not suitable if outcome is unpromising.

Table 2: Contraindications for extracorporeal membrane oxygenation therapy.

End-stage disease
Untreatable underlying disease and congenital malformations
Significant neurological impairment, genetic abnormalities (e.g., trisomy 13 and 18)
Severe, irreversible organ dysfunction
Extreme prematurity (gestational age <35 weeks)
Severe coagulopathy or contraindication for anticoagulation

Neck cannulation has the advantage of allowing an expeditious procedure in an emergency situation, even during mechanical cardiopulmonary resuscitation without interrupting chest compression. A transverse skin incision is made over the lower third of the sternocleidomastoid on the right side. The internal jugular vein, the common carotid artery and the vagal nerve are identified. Oval purse-string sutures are placed in



Figure 1: The ECMO circuit used at the Children's hospital Zurich including a Thoratec Centrimag[®] pump, a heater unit, a Sechrist Air-Oxygen-Mixer and a Polymethylpentene hollow-fibre oxygenator.

vein (VV-ECMO) and artery (VA-ECMO). Selected cannulas, according to the weight of the patient and desired achievable flow rate, are inserted through a vertical incision within the purse-string suture, tightened over tourniquets and connected to the ECMO circuit.

A heparin bolus of 50–100 IU/kg is administered independently of the cannulation site prior to inserting the cannulas. Subsequently, continuous heparin infusion is titrated using a goal activated clotting time in the range of 160–180 seconds. The platelet count is maintained at or above 100 000/mm³. One perioperative injection of prophylactic antibiotic (cefazolin) is administered at each ECMO implantation.

The role of echocardiography for paediatric patients supported with ECMO

Echocardiography should be used preoperatively for patient selection, perioperatively for cannula placement and filling of the left ventricle and postoperatively for surveillance, optimisation, troubleshooting and evaluation of myocardial recovery. Whenever possible, echocardiography is performed before ECMO cannulation to confirm significant systolic ventricular dysfunction as indication for implantation and to

eliminate residual haemodynamic lesions that could be resolved other than by ECMO installation. Furthermore, echocardiography is used to rule out cardiac anomalies, and assure that there are competent valves and normal systemic veins.

During ECMO implantation, echocardiography can guide the insertion and correct placement of the cannulas and document the volume state of the left ventricle. If VV-ECMO with a dual-lumen cannula (AVALON®, Avalon Laboratories, LLC, California, USA) is inserted via a single site (internal jugular vein), we use echocardiographic or x-ray guidance in the cardiac catheter laboratory to ensure adequate orientation of the ports of the cannula (fig. 2). Their proper position must be confirmed with echocardiography, especially the location of the outflow lumen, which must be positioned in the centre of the right atrium with flow directed towards the tricuspid valve. The hepatic veins and the distal inferior vena cava should not be congested.

The first echocardiogram after ECMO implantation on the intensive care unit (ICU) verifies again the localisation of the cannulas. Complications arising from the cannulas include displacement, malposition or obstruction [7].

Repeated echocardiographic monitoring in a child on ECMO is mandatory and includes the course of left ventricular size and contractility, aortic valve opening and mitral valve regurgitation. If the aortic valve is not opening and there is no shunt, the left ventricle of a patient on VA-ECMO will not unload. If the left ventricle than dilates, recovery is not possible. In this case an additional cannula (vent) has to be implanted in the left

ventricle or atrium, or an atrioseptostomy is needed, to ensure sufficient unloading.

Complications must be anticipated and specifically screened for, and are often associated with either the cannula itself or the anticoagulation regimen. Anticoagulation can lead to bleeding, resulting in pericardial effusion and subsequently tamponade, or formation of a thrombus either in the heart, the vessels or the cannulas.

Patient management on the ICU

In order to achieve the desired goal of optimal pulmonary and/or myocardial support and with optimal balance of blood flow, oxygen delivery and oxygen consumption, the following aims are crucial:

1. Provision of adequate oxygenation and CO₂ removal in pulmonary dysfunction.
2. Provision of adequate blood flow to match metabolic needs in patients with insufficient cardiac output.
3. Prevention of complications from other therapies as well as from the ECMO therapy itself.

Numerous considerations need to be addressed when caring for ECMO patients, as they will influence course and outcome (e.g., anticoagulation, nutrition, infection, renal function, psychosocial family support and ethical considerations). Nevertheless, it is important at the beginning of an ECMO run to have a clear view of the likely length of support required and the ultimate destination. This may be complete recovery, bridge to decision, bridge to transplant or bridge to longstanding cardiac support such as a ventricular assist device.

ECMO support might only be necessary for as little as 3 to 6 days (e.g., myocardial recovery after intraoperative stun, persistent pulmonary hypertension of the newborn or vasoplegic shock in sepsis) [8], up to 10 to 14 days, or even significantly longer (e.g., adult respiratory distress syndrome, pneumonia, meconium aspiration syndrome, myocarditis) [9–11]. Time on ECMO support should be used wisely if uncertainties about the underlying disease process remain. If echocardiography leaves questions unanswered, a cardiac catheter study can usually safely be undertaken and should meticulously investigate the patient's haemodynamic state or search for residual lesions, as this will probably have direct implications for the ongoing ECMO support and for immediate decisions [12]. The patient on ECMO needs precise and fastidious anticoagulation management, because bleeding, haemolysis and thrombosis should be prevented as far as possible, as they are primarily responsible for neurological sequelae and dismal outcome [13]. Near-infrared spectroscopy (NIRS) has




Figure 2: Implanted VV-ECMO using a single-lumen cannula (AVALON) showing proper placement in the catheter laboratory.

evolved as a reliable tool to guide therapy in order to improve neurological outcomes after ECMO therapy [14]. NIRS provides indirect monitoring of venous drainage and arterial perfusion. In contrast to echocardiography, NIRS allows permanent monitoring [15].

Nursing care

The monitoring and care of children with ECMO therapy is a special nursing challenge and requires close multidisciplinary cooperation. The ICU nurses are specifically trained for ECMO therapy and are therefore able to respond to acute changes at any given time [17, 18]. During ECMO support children are rest-ventilated, sometimes with an open sternum due to a delayed sternal closure after central cannulation. Any manipulation of the positioning of the patient (e.g., wound care, pressure ulcer care or changing endotracheal tube straps) may change the position of the ECMO cannulas, which can negatively affect the ECMO flow and may even lead to failure of the ECMO circuit. Therefore, ECMO specialist nurses need to be extremely vigilant and respond to these changes. The risks of infections, cerebral haemorrhage and renal failure (e.g., due to decreased renal perfusion during non-pulsatile ECMO flow) are increased and will rise with increasing duration of ECMO therapy. Therefore, careful clinical monitoring of the patient, as well as meticulous antiseptic precautions at the cannulation site, are vitally necessary. ECMO patients are exposed to an increased risk of bleeding [19]. Because of bleeding and blood trauma, some patients will need to have blood products several times daily [20]. The nurses continuously monitor every organ system and have to be able to independently and competently respond to emergencies at any time. Ideally, there are always two nurses responsible for one child on ECMO, sharing the complex supervision and care of the patient. Our patients (children and adolescents) are always part of their personal environment – their parents and other family members. Initially, the parents and families of the children experience a tremendous shock. Especially with neonatal patients, building a relationship with their newborn child is a big challenge for parents. They are not able to hold and care for their child [21], and the uncertainty of the outcome aggravates the situation. It is part of the nursing care to support relatives during this time, as well as during grieving if the outcome is not favourable. The regular presence of parents and family at the bedside is very important for a critically ill child. To support and to build a solid relationship, as well individual contact with their child, it is important to integrate the parents in nursing inter-

ventions such as oral care, skin care, changing nappies or massages. A soothing touch, the parents’ and siblings’ voices, personal toys, music and rituals can be very helpful for the ill child, and also for the family. Parents and family members often need detailed information while their child is on ECMO support and, depending on their personalities, are also often themselves in need of encouragement and personal support, a task the ECMO specialist nurse should provide. Nurses are in a particular situation to care for these highly complex patients, and to support the whole family (table 3).

Table 3: Responsibilities of the nursing team caring for children on ECMO.

Responsibilities of the ECMO nurse	
	Psychological family support
	Visual inspection of the ECMO circuit
	Regulation of ECMO flow, sweep gas and oxygenation
	Anticoagulation management
	Management of the heparin infusion and administration of blood products
	Haemofiltration
	Emergency measures and troubleshooting of the ECMO circuit until perfusionists take over
Responsibilities of the bedside nurse	
	Basic care
	Intravenous fluid administration and medical therapy
	Feeding and administration of parenteral nutrition
	Vital signs monitoring
	Monitoring of fluid intake and output
	Nursing documentation
	Nursing assessment, nursing process, nursing diagnoses

Conclusion

Over the last decades ECMO has become the most frequently used form of extracorporeal life support in children. Depending the underlying disease, it can be modified for venovenous blood flow, to provide sufficient oxygenation of the blood in isolated respiratory failure, or for venoarterial support in the case of cardiac and/or respiratory failure. Echocardiography plays an important role at every step of ECMO, starting with the diagnosis, and including documentation of cannula position, and guidance during surgical cannulation and during troubleshooting. In contrast to adult patients, neck cannulation or central cannulation via median sternotomy are the preferred sites in children. Special circuits are available for the paediatric population in order to keep priming volume low, guarantee optimal flow and oxygenation, and reduce complica-

tion rates. Special management on the ICU is crucial to prevent or recognise complications. Interaction between the various disciplines involved is indispensable to identifying the ideal timeframe for successful weaning. Involvement of the parents throughout all stages of therapy is important and may have a positive impact.

Disclosure statement

No financial support and no other potential conflict of interest relevant to this article was reported.

References

- Guerguerian AM, Ogino MT, Dalton HJ, Shekerdemian LS. Setup and maintenance of extracorporeal life support programs. *Pediatr Crit Care Med*. 2013;14(5, Suppl 1):S84–93. <http://dx.doi.org/10.1097/PCC.0b013e318292e528>
- Barbaro RP, Odetola FO, Kidwell KM, Paden ML, Bartlett RH, Davis MM, et al.; Analysis of the Extracorporeal Life Support Organization Registry. Association of hospital-level volume of extracorporeal membrane oxygenation cases and mortality. Analysis of the extracorporeal life support organization registry. *Am J Respir Crit Care Med*. 2015;191(8):894–901. <http://dx.doi.org/10.1164/rccm.201409-1634OC>
- Freeman CL, Bennett TD, Casper TC, Larsen GY, Hubbard A, Wilkes J, et al. Pediatric and neonatal extracorporeal membrane oxygenation: does center volume impact mortality? *Crit Care Med*. 2014;42(3):512–9. <http://dx.doi.org/10.1097/01.ccm.0000435674.83682.96>
- Domico MB, Ridout DA, Bronicki R, Anas NG, Cleary JP, Cappon J, et al. The impact of mechanical ventilation time before initiation of extracorporeal life support on survival in pediatric respiratory failure: a review of the Extracorporeal Life Support Registry. *Pediatr Crit Care Med*. 2012;13(1):16–21. <http://dx.doi.org/10.1097/PCC.0b013e3182192c66>
- Kurkluoglu M, Hynes CF, Alfares FA, El-Sayed Ahmed MM, Peer SM, Zurakowski D, et al. Choice of Peripheral Venoarterial Extra-Corporeal Membrane Oxygenation Cannulation Site in Patients Above 15 kilograms. *J Card Surg*. 2015;30(5):461–5. <http://dx.doi.org/10.1111/jocs.12538>
- Field ML, Al-Alao B, Mediratta N, Sosnowski A. Open and closed chest extrathoracic cannulation for cardiopulmonary bypass and extracorporeal life support: methods, indications, and outcomes. *Postgrad Med J*. 2006;82(967):323–31. <http://dx.doi.org/10.1136/pgmj.2005.037929>
- Catena E, Tasca G. Role of echocardiography in the perioperative management of mechanical circulatory assistance. *Best Pract Res Clin Anaesthesiol*. 2012;26(2):199–216. <http://dx.doi.org/10.1016/j.bpa.2012.02.005>
- MacLaren G, Butt W, Best D, Donath S. Central extracorporeal membrane oxygenation for refractory pediatric septic shock. *Pediatr Crit Care Med*. 2011;12(2):133–6. <http://dx.doi.org/10.1097/PCC.0b013e3181e2a4a1>
- Rajagopal SK, Almond CS, Laussen PC, Rycus PT, Wypij D, Thiagarajan RR. Extracorporeal membrane oxygenation for the support of infants, children, and young adults with acute myocarditis: a review of the Extracorporeal Life Support Organization registry. *Crit Care Med*. 2010;38(2):382–7. <http://dx.doi.org/10.1097/CCM.0b013e3181bc8293>
- Prodhan P, Stroud M, El-Hassan N, Peeples S, Rycus P, Brogan TV, et al. Prolonged extracorporeal membrane oxygenator support among neonates with acute respiratory failure: a review of the Extracorporeal Life Support Organization registry. *ASAIO J*. 2014;60(1):63–9. <http://dx.doi.org/10.1097/MAT.0000000000000006>
- Smalley N, MacLaren G, Best D, Paul E, Butt W. Outcomes in children with refractory pneumonia supported with extracorporeal membrane oxygenation. *Intensive Care Med*. 2012;38(6):1001–7. <http://dx.doi.org/10.1007/s00134-012-2581-5>
- Panda BR, Alphonso N, Govindasamy M, Anderson B, Stocker C, Karl TR. Cardiac catheter procedures during extracorporeal life support: a risk-benefit analysis. *World J Pediatr Congenit Heart Surg*. 2014;5(1):31–7. <http://dx.doi.org/10.1177/2150135113505297>
- Dalton HJ, Garcia-Filion P, Holubkov R, Moler FW, Shanley T, Heide-mann S, et al.; Eunice Kennedy Shriver National Institute of Child Health and Human Development Collaborative Pediatric Critical Care Research Network. Association of bleeding and thrombosis with outcome in extracorporeal life support. *Pediatr Crit Care Med*. 2015;16(2):167–74. <http://dx.doi.org/10.1097/PCC.0000000000000317>
- de Mol AC, Liem KD, van Heijst AF. Cerebral aspects of neonatal extracorporeal membrane oxygenation: a review. *Neonatology*. 2013;104(2):95–103. <http://dx.doi.org/10.1159/000351033>
- Hoffman GM, Brosig CL, Mussatto KA, Tweddell JS, Ghanayem NS. Perioperative cerebral oxygen saturation in neonates with hypoplastic left heart syndrome and childhood neurodevelopmental outcome. *J Thorac Cardiovasc Surg*. 2013;146(5):1153–64. <http://dx.doi.org/10.1016/j.jtcvs.2012.12.060>
- Butt W, Heard M, Peek GJ. Clinical management of the extracorporeal membrane oxygenation circuit. *Pediatr Crit Care Med*. 2013;14(5, Suppl 1):S13–9. <http://dx.doi.org/10.1097/PCC.0b013e318292ddc8>
- Kotani Y, Honjo O, Davey L, Chetan D, Guerguerian AM, Gruenwald C. Evolution of technology, establishment of program, and clinical outcomes in pediatric extracorporeal membrane oxygenation: the “sickkids” experience. *Artif Organs*. 2013;37(1):21–8. <http://dx.doi.org/10.1111/aor.12032>
- Anderson JM, Boyle KB, Murphy AA, Yaeger KA, LeFlore J, Halamek LP. Simulating extracorporeal membrane oxygenation emergencies to improve human performance. Part I: methodologic and technologic innovations. *Simul Healthc*. 2006;1(4):220–7. <http://dx.doi.org/10.1097/01.SIH.0000243550.24391.ce>
- Larsen R. *Anästhesie und Intensivmedizin in Herz-, Thorax- und Gefässchirurgie*. 9th ed. Heidelberg: Springer Verlag; 2005: p. 2.
- Jöhling B, et al. *Pflege bei Kindern an ECMO oder Kunsttherz*. Z Geburtshilfe Neonatol. 2006.
- Lewis AR, Wray J, O’Callaghan M, Wroe AL. Parental symptoms of posttraumatic stress after pediatric extracorporeal membrane oxygenation*. *Pediatr Crit Care Med*. 2014;15(2):e80–8. <http://dx.doi.org/10.1097/PCC.0000000000000036>

Correspondence:

Martin Schweiger
MD, Priv.Do.
Division of Congenital
Cardiovascular Surgery
University Children’s
Hospital
Steinwiesenstrasse 75
CH-8032 Zurich
[martin.schweiger\[at\]
kispi.uzh.ch](mailto:martin.schweiger[at]kispi.uzh.ch)